Table 7B.7: Tumors of the Musculoskeletal System, by Age, United States

| | | Age Distribution [1] (%), in Years, by Site, 2010-2014 | | | | | | | | |
|-------------------------|---------------------|--|------------------|--------------|--------------|--------------|--------------|------------|-----------|--|
| | < 20 | 20-34 | <u>35-44</u> | <u>45-54</u> | <u>55-64</u> | <u>65-74</u> | <u>75-84</u> | <u>85+</u> | All Cases | |
| Diagnosis | | | | | | | | | | |
| Bones and Joints | 26.4% | 15.7% | 9.5% | 11.8% | 13.1% | 11.7% | 8.0% | 3.7% | 4,126 | |
| Myeloma | 0.0% | 0.6% | 2.9% | 10.8% | 23.3% | 29.2% | 24.1% | 9.2% | 30,340 | |
| Soft Tissue | 8.3% | 9.0% | 8.8% | 14.6% | 19.2% | 17.7% | 14.8% | 7.5% | 15,454 | |
| Death | | | | | | | | | | |
| Bones and Joints | 12.2% | 14.5% | 5.6% | 9.7% | 13.3% | 15.5% | 16.4% | 12.8% | 7,203 | |
| Myeloma | 0.0% | 0.1% | 0.9% | 5.2% | 15.9% | 27.0% | 32.3% | 18.6% | 58,167 | |
| Soft Tissue | 3.5% | 6.1% | 6.0% | 12.8% | 19.7% | 20.8% | 19.2% | 11.9% | 22,683 | |
| Prevalence of Childhood | [2] Bone Cancers by | Age, United Sta | ites, January 1, | 2014 | | | | | | |
| | _ | 0-19 | 20-29 | 30-39 | 40-49 | 50-59 | 60 & Over | All Ages | | |
| Osteosarcoma [3] | Number | 2,048 | 2,585 | 1,772 | 1,440 | 1,323 | 1,077 | 10,245 | | |
| | % of Total | 20% | 25% | 17% | 14% | 13% | 11% | | | |
| Ewing's Sarcoma [4] | Number | 1,487 | 1,482 | 1,115 | 692 | 450 | 258 | 5,484 | | |
| | % of Total | 27% | 27% | 20% | 13% | 8% | 5% | | | |

^[1] Age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130).

Source: Howlader N, Noone AM, Krapcho M, Miller D, Bishop K, Kosary CL, Yu M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (eds). SEER Cancer Statistics Review, 1975-2014, National Cancer Institute. Bethesda, MD, https://seer.cancer.gov/csr/1975 2014/, based on November 2016 SEER data submission, posted to the SEER web site, April 2017. Table 1.11: Age Distribution of Diagnosis by Site, 2010-2014, All races, Both Sexes and Table 1.13: Age Distribution of Deaths by Site, 2010-2014, All races, Both Sexes. Accessed January 22, 2018.

^[2] Number of people diagnosed with cancer as children (ages 0-19) in the United States and alive on January 1, 2014.

^[3] The most common type of primary bone cancer is found in the matrix that forms normal bones. Most osteosarcomas occur in children and young adults in areas where bone is growing rapidly. The most common sites for tumors are the arms, legs, and pelvis.

^[4] Most Ewing tumors occur in bones, with the most common sites being the pelvis, chest wall (ribs or shoulder blades), and the legs, mainly the middle of the long bones. Most Ewing tumors occur in children and teens.