

Table 2B.5.0.1: Hospital Procedures Performed on Persons with Spinal Curvature Diagnosis at Admission, United States, 2010-2014

	Proportion of Total Discharges with Spinal Curvature Diagnosis and Procedure									
	Total Discharges/Visits (in 000s)	Monitoring Deformity	Discectomy	Decompression	Incision/Excision	Fusion	Instrumentation	Disk Replacement	Blood Transfusion	Any Spinal Procedure
Idiopathic scoliosis	161.0	4.1%	6.4%	3.8%	11.9%	17.6%	10.3%	0.2%	10.5%	20.7%
Acquired/secondary scoliosis	20.3	11.9%	19.9%	9.4%	34.0%	47.6%	31.0%	*	17.6%	53.6%
Scoliosis	166.6	3.9%	5.4%	3.5%	10.8%	16.0%	8.8%	0.2%	10.4%	19.4%
Kyphosis	44.9	7.2%	13.9%	7.3%	20.0%	32.0%	22.0%	0.3%	11.8%	35.3%
Lordosis	4.1	12.7%	22.3%	9.4%	32.9%	46.6%	32.4%	*	14.9%	51.2%
Spondylolisthesis	144.6	14.2%	41.0%	17.3%	50.7%	79.2%	57.7%	0.7%	9.8%	84.6%
Sagittal Deformity	190.5	12.4%	34.2%	14.8%	43.0%	67.4%	48.7%	0.6%	10.3%	72.3%
All Spinal Curvature Disorders	357.0	8.4%	20.7%	9.5%	28.0%	43.4%	30.1%	0.4%	10.3%	47.6%

* Estimate does not meet standards for reliability

[1] Source: HCUP National Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2010-2014. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/nisoverview.jsp