Table 2A.17: Hospital Discharges with Back Pain Diagnoses Where Discharge Included Primary (1st) Spinal Fusion Procedure, by

 Demographic Groups, United States 2013

	All Back Pain Diagnoses Discharges				All Primary (1st) Spinal Fusion Discharges [1]					
	<u>Total</u> <u>Diagnoses</u> (in 1,000s)	<u>Average</u> <u>Age</u>	<u>Mean Length</u> of Stay (in days)	<u>Mean</u> Charges	<u>Total</u> Discharges (in 1,000s)	<u>Average</u> Age	<u>Mean Length</u> <u>of Stay</u> (in days)	_(<u>Mean</u> Charges	Proportion Back Pain Discharges with Fusion
Total Discharges	2,977.0	61.0	4.8 \$	52,630	434.0	56.4	3.9	\$ 12	10,960	14.6%
Sex										
Male	1,306.0	59.6	4.9 \$	56,750	201.7	56.3	4.2	\$ 12	14,600	15.4%
Female	1,670.3	62.5	4.7 \$	49,400	232.3	56.5	3.7	\$ 10	07,800	13.9%
Age										
<18	48.3	11.6	6.3 \$	92,020	11.7	13.3	6.3	\$ 18	33,940	24.2%
18-44	487.6	34.1	4.5 \$	49,700	73.0	36.3	3.8	\$ 10	07,850	15.0%
45-64	1,076.0	55.2	4.6 \$	55,190	206.1	55.0	3.5	\$ 10	04,760	19.2%
65 & Over	1,364.9	77.0	5.0 \$	50,250	143.1	72.2	4.3	\$ 12	15,640	10.5%
Race/Ethnicity										
White, Non-Hispanic	2,196.5	62.6	4.7 \$	51,640	329.2	57.3	3.7	\$ 10	07,940	15.0%
Black, Non-Hispanic	297.3	55.5	5.2 \$	51,660	35.5	53.8	5.0	\$ 12	18,330	11.9%
Hispanic	193.1	54.8	5.2 \$	69,180	24.9	52.0	5.0	\$ 1	50,970	12.9%
Other, Non-Hispanic	118.8	58.7	5.3 \$	61,470	18.3	54.0	4.6	\$ 13	30,130	15.4%
Geographic Region										
Northeast	516.4	61.6	5.1 \$	50,450	71.3	55.0	4.1	\$ 9	9,750	13.8%
Midwest	717.3	61.5	4.6 \$	41,320	95.5	56.1	3.7	\$ 9	91,040	13.3%
South	1,142.5	60.4	4.8 \$	50,600	179.1	56.7	3.8	\$ 10	07,760	15.7%
West	600.5	61.3	4.7 \$	73,250	88.1	57.2	4.2	\$ 14	49,320	14.7%

[1] Does not include refusion procedures.

Source: HCUP Nationwide Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2013. Agency for Healthcare Research and Quality, Rockville, MD. <u>www.hcup-us.ahrq.gov/nisoverview.jsp</u>