

Table 2A.20: Primary (1st) Diagnosis for Spine Diskectomy Procedures, United States 2011 and 2013

		2011		2013	
<u>ICD-9-CM</u>			<u>% of Total Spine</u>		<u>% of Total Spine</u>
<u>Diagnosis</u>			<u>Discharges with</u>		<u>Discharges with</u>
<u>Code</u>	<u>Diagnosis</u>	<u>Number</u>	<u>Diskectomy Procedure</u>	<u>Number</u>	<u>Diskectomy Procedure</u>
722.10	Lumbar Disc Displacement	104,389	28.3%	73,785	23.3%
722.00	Cervical Disk Displacement	49,892	13.5%	37,920	12.0%
722.52	Lumbar Disc Degeneration	33,283	9.0%	27,150	8.6%
724.02/.03	Lumbar Spinal Stenosis wo/w neurogenic claudication	22,755	6.2%	23,120	7.3%
721.00	Cervical Spondylosis	21,146	5.7%	18,820	5.9%
722.71	Cervical Disk Disorder	19,310	5.2%	17,745	5.6%
721.10	Cervical Spondylosis with Myelopathy	18,562	5.0%	20,465	6.5%
738.40	Acquired Spondylolisthesis	17,098	4.6%	18,175	5.7%
721.30	Lumbosacral Spondylosis	13,259	3.6%	13,600	4.3%
722.40	Cervical Disc Degeneration	12,546	3.4%	10,640	3.4%
723.00	Cervical Spinal Stenosis	11,052	3.0%	10,745	3.4%
756.12	Spondylolisthesis	5,979	1.6%	6,840	2.2%
722.73	Thoracic Disc Disorder with Myelopathy	3,593	1.0%	3,390	1.1%
996.49	Complication of Internal Orthopaedic Device			3,540	1.1%
737.30	Idiopathic Scoliosis			2,530	0.8%
722.83	Postlaminectomy syndrome, lumbar region			2,200	0.7%
724.40	Thoracic or lumbosacral neuritis or radiculitis, unspecified			1,485	0.5%
723.40	Brachial neuritis or radiculitis NOS (cervical region)			1,395	0.4%
737.39	Other kyphoscoliosis and scoliosis			1,200	0.4%
	All Other diagnoses	36,602	9.9%	21,940	6.9%
	All Primary (1 st) Diagnosis for Diskectomy Procedure	369,466	100.0%	316,685	100.0%

Source: HCUP Nationwide Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2011, 2013. Agency for Healthcare Research and Quality, Rockville, MD. www.hcup-us.ahrq.gov/nisoverview.jsp