

**Table 3A.4.1.2: Bed and Lost Work Days Associated with Self-Reported Doctor-Diagnosed Arthritis and With Any Medical Conditions for Adults Age 18 and Older, by Age, United States 2013-2015**

	Age			Total
	18 to 44 Years	45 to 64 Years	65 Years & Older	
<b>Adults With Self-Reported Doctor-Diagnosed Arthritis [1]</b>				
Adults with Reported Condition [1] (in 000s)	7,965.5	24,181.0	22,226.0	54,372.5
% Total Population [3]	6.9%	29.1%	49.7%	21.9%
<b>Bed Days</b>				
Adults with Bed Days [4] (in 000s)	4,666.2	12,093.1	7,845.5	24,604.8
% of Total Population	4.1%	14.6%	17.5%	9.9%
% of Population with DDA	58.6%	50.0%	35.3%	45.3%
Average Number Bed Days in Past 12 Months	20.1	26.4	24.7	24.7
Total Bed Days (in millions)	93.8	319.1	194.0	607.0
Share of Total Bed Days by Group	15%	53%	32%	
<b>Lost Work Days</b>				
Adults with Lost Work Days [5] (in 000s)	3,711.8	7,658.3	1,270.0	12,640.0
% of Total Population	3.2%	9.2%	2.8%	5.1%
% of Population with DDA	46.6%	31.7%	5.7%	23.2%
Average Number Work Days Lost in Past 12 Months	13.4	14.6	15.3	14.3
Total Lost Work Days (in millions)	49.9	111.6	19.4	180.9
Share of Total Lost Work Days by Group	28%	62%	11%	
<b>Adults Reporting Any Medical Condition [2]</b>				
Adults with Reported Condition [1] (in 000s)	73,622.5	69,684.2	42,423.6	185,730.4
% Total Population [3]	64.2%	83.9%	94.9%	75.0%
<b>Bed Days</b>				
Adults with Bed Days [4] (in 000s)	34,082.0	28,697.9	13,136.6	75,916.5
% of Total Population	29.7%	34.5%	29.4%	30.6%
% of Population with Any Medical Condition	46.3%	41.2%	31.0%	40.9%
Average Number Bed Days in Past 12 Months	9.3	17.3	22.0	14.5
Total Bed Days (in millions)	316.6	496.8	289.1	1,102.3
Share of Total Bed Days by Group	29%	45%	26%	
<b>Lost Work Days</b>				
Adults with Lost Work Days [5] (in 000s)	31,239.0	22,926.4	2,321.1	56,486.4
% of Total Population	27.2%	27.6%	5.2%	22.8%
% of Population with Any Medical Condition	42.4%	32.9%	5.5%	30.4%
Average Number Work Days Lost in Past 12 Months	8.2	10.8	12.6	9.4
Total Lost Work Days (in millions)	255.2	248.5	29.2	533.2
Share of Total Lost Work Days by Group	48%	47%	5%	
<b>Proportion Attributed to Arthritis versus Any Medical Condition</b>				
Adults with DDA	10.8%	34.7%	52.4%	29.3%
Adults with Bed Days	13.7%	42.1%	59.7%	32.4%
Total Bed Days	29.6%	64.2%	67.1%	55.1%
Adults with Lost Work Days	11.9%	33.4%	54.7%	22.4%
Total Lost Work Days	19.5%	44.9%	66.4%	33.9%

[1] Responded "yes" when asked: Have you ever been told by a doctor or other health professional that you have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia?

[2] Responded "yes" when asked if have a medical condition, including circulatory (coronary, heart), respiratory, musculoskeletal, hearing trouble, seeing trouble, emotional disorder, mental health disorder, diabetes, cancer, recurring severe headaches, cognitive function issue, or sleeping issue. Respondents may indicate more than one medical condition.

[3] Adjusted to working age population aged 18 and older.

[4] A bed day is defined as 1/2 or more days in bed due to injury or illness in past 12 months, excluding hospitalization.

[5] A missed work day is defined as absence from work due to illness or injury in the past 12 months, excluding maternity or family leave.

Source: National Health Interview Survey (NHIS)\_Adult sample, 2013, 2014, 2015. <http://www.cdc.gov/nchs/nhis/data-questionnaires-documentation.htm> July 23, 2016.